

Phage Lambda DNA/*Hind* III Markers.

Description Lambda DNA (cl857 *Sam 7*)/*Hind* III Markers are prepared by digesting Lambda DNA with *Hind* III, followed by heat-inactivation of the enzyme. The DNA fragments are then ethanol-precipitated and resuspended in storage buffer.

Fragments (Base Pairs) 1- 23,130; 2- 9,4; 3- 6,557; 4- 4,361; 5- 2,322; 6- 2,027; 7- 564, 8 - 125

Storage Buffer 10mM Tris-HCl (pH 8.0), 1mM EDTA.


Concentration: 0.2-0.5µg/µl.

Number of Bands: 8.

Storage Store at -20°C.

Loading The recommended amount of size marker to load on an agarose gel is 1µg/lane.

Attention! The cohesive ends of fragments 1 and 4 may cause formation of extra band 27491 bp. The fragments may be separated by heating to 65°C for 3 minutes before loading the sample on the gel.



<u>Cat#</u>	<u>Pack size</u>	<u>Price, Euro</u>
302005	0,05mg	
302025	0,25mg	

Version 1.1 01.03

Bioron GmbH

Contact: Phone: +49-(0)-621- 5720 915 Fax:+49-(0)-621-5720 916

E-Mail: info@bioron.net NET: www.bioron.net