



The ENZYME Company

Enzyme	Prototype	Recognition Sequence	Cat No	Package, u.a.
--------	-----------	----------------------	--------	---------------

PspE I	BstE II	G [^] GTNACC CCANTG [^] G	238020	2000 units
			238100	10000 units

Lot-number:	Assayed:	Quantity:

Origin	Pseudomonas species E
Concentration	5000-10000, u.a./ml
Storage conditions	10 mM Tris-HCl (pH 7.5); 50 mM KCl; 0,1 mM EDTA; 7 mM 2-mercaptoethanol; 200 ug/ml BSA; 50% glycerol; Store at -20°C.
Ligation	After 5-fold overdigestion with enzyme more than 90% of the DNA fragments can be ligated and recut.
Non-specific activity	No nonspecific activity was detected after incubation of 1 ug of DNA with 10 u.a. of enzyme for 16 hours at 37°C.
Optimum temperature	37 °C
Inactivation 20 minutes under 65 °C	Yes
Optimum SE-buffer	B (10 mM Tris-HCl (pH 7.6 at 25°C); 10 mM MgCl ₂ ; 1 mM DTT.)

Enzyme activity in % of maximum :

B	G	O	W	Y
100	50 - 75	25 - 50	50 - 75	50 - 75

References: Dedkov, V.S., Rechkunova, N.I., Shevchenko, A.V., Degtyarev, S.Kh. Unpublished observations (1994).

Unit-definition	One unit of the enzyme is the amount required to hydrolyze 1 µg of DNA in 1 hour in a total reaction volume of 50 µl. Concentrated enzymes are diluted to approximately 1000 units/ml with the buffer (10mM Tris-HCL (ph7.6); 50 mM KCL; 0,1 mM EDTA; 1 mM DTT; 200 µg/ml BSA; 50% glycerol) before determining their activity.
------------------------	---

Bioron GmbH

Contact: Phone: +49-(0)-621- 5720 915 Fax:+49-(0)-621-5720 916
E-Mail: info@bioron.net NET: www.bioron.net